

# The Phytotoxic Virulence Factor BcSnod1 from *Botrytis cinerea*

Gnana Sreekar Nidadavolu, Hana McFeeters, Robert McFeeters  
Department of Chemistry

## Overview

*Botrytis cinerea* is a necrotrophic fungal pathogen known to infect over a 230 species of plants across the globe. It secretes a 12.4 kDa pathogenic virulence factor BcSnod1 which belongs to a new class of phytopathogenic proteins termed ceratoplatanins. Ceratoplatanins are known to stimulate plant defense systems and develop hypersensitive responses in hosts that ultimately lead to cell death. We aim to understand the properties of BcSnod1 and identify the molecular mechanisms of its effect on plants. We have developed a highly efficient recombinant yeast expression system for BcSnod1 and purify it by ultra-filtration and size exclusion chromatography. The resulting protein is used for testing against tomato and tobacco plants.

## Impact

*B. cinerea* infections occur across the globe and result in 20% of overall crop losses worldwide. In the US alone, crop losses to *B. cinerea* are estimated at 2 billion USD per year with another 15-20 million spent on anti-botrytis products each year. With emergence of fungal drug resistance and increasing demand for global food production, it is important to limit agricultural losses to *Botrytis cinerea*. Considering the wide population of crop plants *B. cinerea* is capable of infecting, cutting edge research is necessary to stop this fungal pathogen. We continue to elucidate the mechanism of pathogenicity, identify phytoalexin responses, and develop transgenic varieties of plants resistant to *B. cinerea* infection. Better understanding of *B. cinerea* infection and BcSnod1 provides new avenues for inhibition from seed to store.

### B. cinerea effect on some common crops



Strawberry



Bell pepper



Grapes



Bulb flower

## Acknowledgements

I thank all my lab members for their help and cooperation.

## Key Findings

- mg quantities of recombinant BcSnod1 have been produced in *P. pastoris*
- Recombinant BcSnod1 is active
- $\mu\text{g}$  quantities of protein cause plant cell death on contact
- The BcSnod1 effect is visible within days
- BcSnod1 is a major necrotrophic factor for *B. cinerea* infection

### Necrosis Timescale



Day 5

Day 7

Day 9

Day 14

Day 18

Injection: 100  $\mu\text{g}$  of BcSnod1 in 20  $\mu\text{l}$  buffer

BcSnod1 was infiltrated into the mid-veins of two-month old tomato and tobacco leaves. Necrosis usually started in 4-5 days post infiltration and eventually spread to the entire leaf. However, in tobacco leaves, infection did not spread to whole leaf but to surrounding areas from the point of infiltration.

### Tobacco leaves infection



Buffer ctrl

BcSnod1 Day7

Injection: 320  $\mu\text{g}$  of BcSnod1 in 100  $\mu\text{l}$  buffer

## Explanation

The project uses recombinant DNA technology to express the fungal virulence factor BcSnod1 in *P. pastoris*. Identification of the interactions leading to the necrotic behavior on the above shown leaves is a driving step for understanding the plant defense responses and phytoalexins production. We further understanding of BcSnod1 and development of transgenic plants more resistant to *B. cinerea* infection.